

RESEARCH

Open Access



Assessing young adults' menopause knowledge to increase understanding of symptoms and help improve quality of life for women going through menopause; a student survey

Vaishvi Patel¹, Sue Ross¹ and Beate C. Sydora^{1*}

Abstract

Background Due to menopause being a largely invisible and under-discussed topic in wider society, women often deal with menopause-related complications on their own. Social support and awareness have been shown to reduce negative menopausal experiences; however, lack of menopause knowledge, particularly among younger people, may deter support for women suffering from menopause symptoms. This study aims to assess the level of knowledge young adults have on menopause to be able to create interventions that target knowledge gaps and increase understanding of women's experiences and difficulties during their menopause transition.

Methods We created an electronic questionnaire based on menopause literature and guidelines from Menopause Societies. It was pilot-tested on young people in the target group age ($n = 14$; 7 male and 7 female), menopause clinicians ($n = 5$), and women experiencing menopause ($n = 4$). The final survey included questions on participant demographics, general menopause knowledge, and options to support menopause management and was distributed through university student newsletters. Responses over a two week period were collected anonymously. Descriptive statistics were applied to characterize participants, define menopause knowledge, and identify gaps. Chi-squared statistics was used for group comparison, and open questions were analyzed using qualitative content analysis.

Results Survey responses were collected from 828 students; the average age was 22.1 ± 5.1 and 83.6% were female. Participants belonged to all faculties and included students from a variety of family settings and living conditions. Knowledge questions revealed a good understanding of the basic menopause physiology for most respondents, but there were gaps in understanding of symptoms and symptom management. Female sex and personal connection to menopausal women had a positive effect on the degree of menopause knowledge. Both males and females reported increased knowledge confidence at the end of the survey.

Conclusion Our survey provides evidence that young adults of both sexes have a general baseline knowledge of menopause and its symptoms and are open to learning strategies to help support menopausal women. Our findings will assist in developing targeted educational resources to increase social support and awareness, reduce stigma

*Correspondence:

Beate C. Sydora

bsydora@ualberta.ca

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

and improve the quality of life for menopausal women, and help prepare younger women for their future menopause journey.

Keywords Menopause, Menopause symptoms, Menopause experience, Menopause knowledge, Quality of Life, Climacteric symptoms

Background

Menopause is defined as the point in time when a woman has ceased having menstrual periods for 12 consecutive months, marking the end of her reproductive years. During menopause and its transition, estrogen deficiency can give rise to a variety of psychological, physical, and endocrine changes, including mood disorders, sleep disorders, sexual dysfunction, joint pain, body aches, hot flashes, and increased risk for the development of osteoporosis, and cardiovascular health issues [1–4].

Women's perception of menopause severity and symptoms varies widely and can be influenced by individual experiences related to cultural context, mental state, emotional health, and social support [3]. Studies have shown that women undergoing the menopause transition are psychologically more vulnerable and are at high risk for developing depression and mood disorders; these are attributed to several factors such as inadequate social support which is among the most important [5, 6]. Women are often reticent about menopause, causing it to be an unmentioned, largely invisible topic that is seldom discussed in wider society, leaving women to deal with menopause-related changes on their own. The stigma surrounding menopause, associations with mood changes, the shame of ageing, and the taboo about revealing menopausal symptoms, all contribute to hidden distress. Furthermore, the stigma from others' reactions can become internalized, further decreasing quality of life. The result is a hidden public health disorder that can cause suffering for women as well as their family members [7–9].

Based on our research group's previous work, menopausal women often report feeling anxious, frustrated, and alone because they feel their friends and family are not able to understand what they are experiencing and are not able to provide them with the support they need [8, 9]. Social support and education have been shown to reduce negative menopausal experiences and result in a higher quality of life for women during menopause [10, 11]. It has also been shown to reduce the risk of mental health disorders in women during the menopause transition [3]. Together these findings suggest that social support, including physical and emotional empathy, can be a simple, cost-effective, and sustainable way to help women through this transition, mitigating menopausal complications. Intervention programs designed to increase

the awareness and education of menopausal women, as well as their spouses, have shown to be effective but there have not been interventions focused on improving the understanding of younger adults of both sexes about women's experience of menopause [2, 10]. Younger adults often consider menopause to be a topic of humor or fear. Educating young adult family members and relatives to better understand the topic of menopause will enable them to be supportive of menopausal women.

Therefore, this study was designed to assess young adults' basic knowledge about the topic of menopause, including its symptoms and management experience, in order to address their knowledge gaps through educational material. In this way, we hope to break the cycle of misunderstanding and create changes that will encourage more sympathetic responses to women experiencing menopause.

Methods

To approach our research goal, we employed a cross sectional design in form of an electronic survey. The methodology and results of the survey were reported using published recommendations such as Checklist for Reporting of Survey Studies (CROSS) and Checklist for Reporting Results of Internet E-Surveys (CHERRIES) to ensure that this study was conducted in a reliable, reproducible, and transparent manner [12, 13].

The study was approved by the University of Alberta Health Research Ethics Board (Pro00108310).

Design of an electronic survey

Our open survey was designed based on the menopause literature review and guidelines from the International Menopause Society (IMS) and the North American Menopause Society (NAMS) [14, 15]. We also sought inspiration from validated menopause questionnaires such as the menopause symptom rating scale (MSRS) [16] and the Menopause-specific Quality of Life (MENQOL) questionnaire [17–19], as well as a symptom severity questionnaire used to follow the progression of menopause symptoms in women treated at the two Menopause Outpatient Clinics in Edmonton, Alberta [20]. In addition, we considered common concerns raised by menopausal women in recently conducted focus groups at these outpatient clinics [8].

The electronic survey was created using the secure web application Research Electronic Data Capture (REDCap) [21] and included a demographic and menopause knowledge section. Within the demographics section, participant demographics such as age, sex, field of study, relationship with menopausal women and family structure and composition were asked. The menopause knowledge section included knowledge-based and opinion-based questions using multiple choice, sliding scale and short answers. Where applicable a non-response option was included.

Pilot-testing of the survey

The first draft of the survey was pilot-tested to gather feedback on its usefulness, comprehensiveness, and adequateness before it was distributed. The survey was administered to (a) young adults (under 30 years of age) that are not current University of Alberta students (b) menopause clinic health professionals, and (c) menopausal women aged 50 and older. The participants were purposefully sampled and recruited through word-of-mouth of known contacts to the research team. The volunteers who agreed to participate received an e-mail containing the formal study information letter (including the consent process) and a unique URL to access a Google Form where they had an opportunity to review questions and provide feedback.

We received feedback from 23 individuals, 14 young adults (7 males, 7 females), 5 menopause health professionals, and 4 menopausal women aged 50 years and older. The survey draft was modified based on the feedback received. The final survey spanned 6 pages and took less than 20 min to complete.

Data collected from the pilot testing was not used in the analysis.

Participant group and survey distribution

The survey was distributed in June 2020 through email to undergraduate and graduate students, enrolled at the University of Alberta through Undergraduate Digest Newsletter and Graduate Student Communications, respectively. The email contained a prompt along with a link to the survey. To partake in the survey, participants needed to meet the following inclusion criteria: currently enrolled at the University of Alberta and at least 18 years of age. A negative answer to these two questions terminated the survey. No specific exclusion criteria were applied. Completing the survey indicated consent.

While participation in the survey was voluntary, we employed a purposive sampling technique; we chose University of Alberta students as our target group as most students would belong within the younger adult demographic. In addition, this target group was most available,

and conducting a survey was an appropriate method during the COVID-19 pandemic when no face-to-face interactions were permitted.

Data acquisition

Study data was collected anonymously during a time spanning two weeks in June 2020 and stored in REDCap, accessible only to the researchers. Students were only able to partake in the survey once and their responses were saved every time an answer was entered. The participants were not able to alter their previous answers once they have proceeded to the next portion of the survey.

Upon completion of the survey, participants were redirected to another site where they could enter their name and email address to participate in a draw for a \$ 50 (CAD) gift card. There was one random gift card draw for every 50 participants.

Data analysis

For analysis, the data was exported into Microsoft Office Excel and descriptive analysis was applied. Questionnaires that were terminated early were not analyzed. Nine evidence-based knowledge questions were included with multiple choices or yes/no answers. Only one correct answer was available for each item. Each correct answer received one point; wrong answers were counted as 0 points. For each participant the points were summed, with the lowest score being 0 and the highest being 9; a higher score indicating more knowledge about menopause.

For statistical analysis of data comparison based on demographic factors, IBM-SPSS data analytics software version 28 was used. A chi-squared test of independence was performed to examine the relationship between sex and menopause knowledge levels and between having close contact with a woman in menopause and levels of menopause knowledge. Paired t-tests were used to compare means of self-assessed levels of menopause knowledge confidence at the start and at the end of the survey. Independent sample tests were used to compare these means between males and females and between participants who have close contact with a woman in menopause or do not. Qualitative content analysis was utilized to generate themes and summarize knowledge gaps from opinion-based and open-ended questions.

Results

Participant characteristics

Out of 929 returned surveys, 828 completed the demographic section. Of these, 76 were graduate and 752 were undergraduate students. Most participants were enrolled in the Faculty of Science (195, 23.6%), followed by Arts (160, 19.3%), and Engineering (93, 11.2%), which

is representative of the size of the programs. The majority of participants were female (683, 82.5%) and identified as a woman (658, 79.5%). We received responses from students of all ethnic backgrounds with the most being Caucasian (432, 52.2%) followed by South Asian (123, 14.9%) and East Asian (111, 13.4%). Of all participants, 460 (55.6%) were single and 477 (57.7%) lived with their families. Only 8.7% of participants did not have some form of a relationship with someone currently in menopause while 81.6% of participants reported having a close

contact with a woman in menopause. The average age of participants was 22.1 ± 5.1 (mean \pm SD). Demographic details are provided in Additional file 1.

Respondents completed knowledge-based questions and questions on support and help for menopausal women. The complete questions and corresponding answer options can be found in Additional file 2.

Menopause knowledge

An open-ended question “please describe what menopause is” was answered by 785 respondents (Table 1).

The majority (66.8%) described menopause as the time when menstrual periods stop and the end of fertility (48.2%), with a few mentioning that it starts 12 months post last menstrual period (2.2%). About a third (33.7%) acknowledge that this has to do with hormonal changes, and a few (4.5%) named estrogen and progesterone as the hormones involved. About a quarter (23.3%) mentioned that menopause relates to debilitating symptoms with a few mentioning prototypical menopause symptoms such as hot flashes and mood swings. Six admitted that they do not know, are not sure, or have no idea what menopause is.

Table 2 depicts the scores of participants on 9 knowledge-based questions. A chi-squared test of independence showed that there was a significant difference between menopause knowledge and sex (X^2 (1, $N=780$) = 11.9, $p=0.02$); females demonstrated higher knowledge levels than males (Table 2A). There was also a significant difference between menopause knowledge level and whether the participants had close contact with a woman in menopause (X^2 (1, $N=787$) = 15.7, $p=0.003$); participants with close contact to a woman

Table 1 Open answer question: “In a sentence, please describe what menopause is?”

Themes N= 785 responses	Topic ^a	N [%]
Final menstruation	End of periods/menstrual cycles	524 [66.8]
	End of fertility	378 [48.2]
Hormonal changes	Change in hormones	229 [29.2]
	Reduction in Estrogen	35 [4.5]
Menopause timing	One year/12 months post last menstruation	17 [2.2]
Age at menopause	Described as post 40, between 40 and 60, or around age 50	85 [10.8]
Symptoms	General experience symptoms	85 [10.8]
	Hot Flashes	53 [6.8]
	Mental/Mood changes	41 [5.2]
	Sleep disturbance	4 [0.5]
Reduced QOL	Suffering/not feeling well	13 [1.7]
No/insufficient knowledge of menopause	“Don't know” or “not sure”	6 [0.8]

^a more than one answer is possible within topics

Table 2 Level of menopause knowledge

A) Knowledge-based questions (n = 9) by biological sex			
Number of correct answers	Female N = 651 N [%]	Male N = 129 N [%]	Prefer not to answer N [%]
9	191 [29.3]	33 [25.6]	4 [57.1]
8	281 [43.2]	49 [38.0]	2 [28.6]
7	122 [18.7]	23 [17.8]	1 [14.3]
6	24 [3.7]	12 [9.3]	0 [0.0]
5 and less	33 [5.1]	12 [9.3]	0 [0.0]
B) Knowledge-based questions (n = 9) by having close contact with a woman in menopause stage			
Number of correct answers	Close contact N = 648 N [%]	No contact N = 139 N [%]	
9	198 [30.6]	30 [21.6]	
8	278 [42.9]	54 [38.8]	
7	116 [17.9]	30 [21.6]	
6	27 [4.2]	9 [6.5]	
5 and less	29 [4.5]	16 [11.5]	

undergoing menopause scored higher than those that did not (Table 2B).

Knowledge about menopause symptoms

To assess knowledge of menopause symptoms, survey respondents were asked to check off all symptoms from a choice of 9 health issues that they believe are associated with menopause. 754 participants responded to this question. The majority correctly identified major menopause symptoms. Mood changes (728, 96.6%), fatigue (689, 91.4%), and hot flashes (682, 90.5%) were the three most picked answers by participants, while other symptoms such as depression, memory loss, aching joints and dry eyes were chosen by 29 to 70% of participants.

Survey participants were also asked to score on a scale of 1 – 10, how debilitating they think menopause symptoms are on average for women. The scoring was done by 762 participants. On average the participants gave the symptom experience a score of 6 in terms of debilitation (Fig. 1).

Menopause perception and opinion

To assess empathy and compassion towards menopausal women, participants were asked a variety of multiple-choice questions. The majority of participants believed that a lower socioeconomic status contributes to an earlier onset of menopause (62.6%) and that social isolation from friends and family is an ineffective way of managing menopausal symptoms (91.2%). Participants also indicated that providing social and emotional support is the most sustainable way to support a woman undergoing menopause (97.5%) (Table 3).

Participants were asked to rate their confidence and knowledge of menopause at the start and end of the survey on a scale from 1–10. Female participants gave themselves a higher score than their male counterparts. Similarly, those with relationship to a woman in menopause gave themselves a higher score than those without such a relationship. All groups of participants gave themselves a significantly higher score at the end of the survey compared to their pre-survey score (Table 4).

When asked the question “Would you be interested in learning more about menopause?”, most participants (674, 90.3%) indicated that they were interested in learning more about menopause with the majority interested in a website with educational resources (414, 55.5%), followed by a lecture delivered by a content expert (216, 29.0%) and physical printouts and brochures (44, 5.9%).

Discussion

This study investigated university students’ knowledge and perception of menopause. We did not intend to provide education or investigate specific treatment options such as hormonal, non-hormonal drug/herbal and laser therapies. The survey findings suggest students have a basic knowledge of the physiology and the symptoms associated with menopause (Tables 1 and 2). In answering an open-ended question, most students mentioned that menopause was associated with hormonal changes and cessation of menstrual periods (Table 1). A few students also recognized hot flashes and mental/mood changes as major menopause symptoms. These two symptoms, in addition to fatigue, were also among the symptoms correctly chosen from a drop-down list as symptoms




Fig. 1 Rating of debilitation associated with menopause symptoms. Participants responded to the question “On a scale of 1–10, how debilitating do you think menopause symptoms are on average for women?” The number of responses at each level from 1 to 10 is shown. N=762, median: 6, mean ± SD: 6.06 ± 1.67

Table 3 Socioeconomic and empathy questions

Possible answers to questions	N [%]
Can lower socioeconomic status contribute to an earlier onset of menopause?	
N = 763	
Yes	478 [62.6]
No	285 [37.4]
Which of the following is a likely scenario occurring due to menopause?	
N = 747	
Janet forgetting where she put her phone	253 [33.9]
Amira slipping on a patch of ice in her driveway	21 [2.8]
Sumin starting to get oily skin and hair, and acne	459 [61.4]
Dyani having regular menses (periods)	14 [1.9]
How can you help a woman undergoing menopause?	
N = 747	
Tell her to get a grip	2 [0.3]
Tell her to take a break	5 [0.7]
Be kind and supportive	738 [98.8]
Leave her alone with her emotions	2 [0.3]
Which of the following is the most sustainable way to support a woman undergoing menopause? N = 747	
Buy her an AC unit	7 [0.9]
Provide social and emotional support	728 [97.5]
Tell her she will get over it	4 [0.5]
Keep your distance from her	1 [0.1]
Buy her a solo vacation to Paris	7 [0.9]
Which of the following is an ineffective way of managing menopausal symptoms?	
N = 747	
Exercising	21 [2.8]
Journaling every day	13 [1.7]
Social isolation from friends and family	681 [91.2]
Seeking medical help	27 [3.6]
Using a calendar to keep track of events	5 [0.7]

Table 4 Self confidence assessment of menopause knowledge

Respondents	N	Confidence level at start of survey [mean ± SD]	Confidence level at end of survey [mean ± SD]	Paired t-test at start versus end	Independent sample test between groups
All respondents	746	5.05 ± 2.06	5.32 ± 1.88	<i>p</i> < .001	na
Female	618	5.19 ± 1.96	5.43 ± 1.81	<i>p</i> < .001	at start: <i>p</i> < .001
Male	121	4.26 ± 2.40	4.76 ± 2.13	<i>p</i> = .006	at end: <i>p</i> < .001
Not disclosed	7	na	na	na	
Close contact to a woman in menopause	617	5.17 ± 1.99	5.42 ± 1.82	<i>p</i> = < .001	at start: <i>p</i> < .001
No close contact to a woman in menopause	129	4.47 ± 2.30	4.86 ± 2.07	<i>p</i> = .04	at end: <i>p</i> = .002

associated with menopause, indicating that students and young adults are aware of the stereotypical symptoms of menopause but may not be aware that menopause can encompass a much greater variety of symptoms including

depression, memory loss, and dry eyes. This is consistent with other studies which show that even middle aged women and spouses of menopausal persons are not fully aware of the more hidden symptoms and experiences of

menopause [22]. Nevertheless, based on the researcher-made scoring system of the questionnaire, most participants scored 7 or 8 out of 9, once again suggesting that students have a modest knowledge of menopause. Although these scores can be influenced by age, past experiences, and program of study, we found that both sex and close contact with a menopausal woman significantly impacted the scores. Females in comparison to males scored statistically higher and an even bigger difference was seen between participants who have close contact with a woman undergoing menopause compared to those without this contact, with the former scoring higher. These scores overlap with the participants' perceived self-confidence scores on menopause knowledge, with females and those in contact with a menopausal person giving themselves a higher score than their counterparts (Table 4). In all cases, participants report a higher self-confidence score post-questionnaire indicating that thinking about menopause and being confronted with the topic is a learning process and boosts confidence in the topic.

The higher scores of female participants was an expected outcome as menopause is an almost universal stage in women's health, but the increase in the menopause knowledge confidence score and the positive influence on scores through a close relationship to a menopausal person was an interesting and promising finding. This goes to show that individuals can be educated and informed about menopause and other potential conditions and that learning can be based on proximity and without a lived experience, similar to the spouses of those undergoing menopause, ultimately making young adults a teachable target demographic [23].

On a scale of 1 – 10, the participants rated the debilitation brought on by menopause on average as 6/10, indicating that there was a general recognition and awareness of the negative impact of menopause on quality of life (Fig. 1). Although none of the participants are likely experiencing menopause currently, they were able to apply their knowledge of the condition to understand the impact it has on quality of life. A similar result was observed in other scenario-based and awareness questions which highlighted participants' empathy and thoughtfulness (Table 3). Studies have shown that the younger generation has the ability to consider others' beliefs, express concern, and that their empathy leads to building social relationships [24, 25]. The aforementioned reasons, in addition to our findings that over 90% of participants indicated interest in learning more about menopause, make young adults an excellent demographic to target for an intervention.

Menopause, with its many complications, has adverse effects on the quality of life of women and places strain

on their relationships, especially with their partners [9, 26]. A study by Currie and Moger which surveyed women 45 years of age or greater with menopause symptoms and their partners found that the physical and emotional effects of menopause affect women's social life, professional lives, and relationships [27]. These women reported more frequent arguments with family members, feelings of isolation, negative impacts on sex life, missed work and decreased enjoyment in social interactions due to their symptoms. The stigma surrounding menopause further contributes to the decreased quality of life and prevents open communication with family members and friends and even deters women from getting help and advocating for themselves.

One potential solution is increasing social and emotional support for these women, which has been positively correlated with a decrease in the symptoms experienced and increases in health and prosocial behaviors and self-esteem [3]. A study by Shariat-Moghani et al., in which husbands of postmenopausal women received educational programming regarding menopause and its complications, resulted in a decrease in negative menopause-related experiences by their spouses [10]. This study along with others confirms that increasing family awareness and support improves women's experiences and psychological and physical health [11, 26, 28]. Another approach is to educate the pre- and peri-menopausal women themselves as most have limited knowledge about menopause, its symptoms, and management and often are interested to learn more about the topic [29, 30]. Studies have found that educational health interventions can decrease somatic and psychological symptoms in premenopausal and menopausal women; the increased awareness helps them to better understand and cope, making it less likely that women would choose to avoid medical treatment [4, 31].

But the most encompassing approach would be educating young adults who will become menopausal women, or spouses, co-workers, neighbours and close contacts, or physicians of menopausal women. By raising awareness in younger people through interventions and methods they most prefer, such as an open-access website with resources, we can prevent the suffering brought on by menopause in both the woman and her family.

Limitations

There are a few limitations to this study. A sampling bias is introduced as only university students were chosen to participate in the survey. Furthermore, many participants had likely even received biology and medical teaching, or the survey title may have attracted or dissuaded a certain subset of students from participating. We also can not exclude the possibility that some students were

attracted to the survey because of the opportunity to win a \$ 50.00 (CAD) gift card and were lesser interested in the topic. Our sample is not representative of the general younger adult population as it overestimates the level of menopause knowledge of the average young adult, making the finding difficult to extrapolate. Nevertheless, it serves as a first step in focusing on this population and assessing their knowledge level. The participants also did not have the option to express their lack of knowledge or uncertainty regarding the survey questions which may lead to answers being guessed, nevertheless, it provides an opportunity for participants to extrapolate from their base knowledge and potentially realize their shortcomings.

Strength of the study

This study has several strengths. It is the first study of its kind, assessing menopause knowledge in younger adults including all sexes and genders. Although a researcher-made survey was used, it was evidence-based and pilot-tested, and feedback was incorporated from field experts and young adults of both sexes to ensure that the questions were applicable to our target demographic. There was also a large sample size (828 participants) including students from all faculties and backgrounds which allows for a closer approximation of knowledge levels in younger adults receiving post-secondary education. Finally, participants on average reported an increased knowledge confidence post-survey, implying that completion of the survey increases awareness about menopause.

Knowledge translation

The result from this study provides an opportunity for further studies to examine the younger adult population to truly understand their knowledge of menopause and other women's health issues. In addition, based on the results we will create free, easy to access, digital resources on menopause and its experience to proactively educate the younger adult population to reduce the stigma around women's health. These resources will arm women and their family members with the knowledge to facilitate intergenerational communication and techniques to support their loved ones, while improving younger women's experience when they reach menopause, and to ultimately create happier and healthier communities.

Conclusion

Symptoms such as hot flashes, night sweats, sleep disturbances and vaginal dryness begin during the menopausal transition and persist in post-menopause and for the rest of some women's lives [32]. These somatic symptoms often present alongside psychological symptoms such as anxiety, depression, and mood fluctuations, reducing

the quality of life for these women [33]. In addition, during the menopause transition, women often experience other changes within their family, social and work life which can bring about other stressors. The amalgamation of these changes and the increasing life expectancy of women emphasize the importance of caring for menopausal women who are experiencing a turning point in their lives.

Findings of this study have highlighted that young adults in an academic setting have a general knowledge of menopause and its complications and possess the ability to empathize with and understand menopausal women. Their knowledge level is cursory and is positively affected by their sex (female) and having a connection to menopausal women. Most participants indicated that they were interested in learning more about menopause through electronic resources. According to these findings, young adults are an excellent demographic for an educational intervention as they demonstrate empathy, teachability and an interest in learning more, therefore if we equip them with the right resources and strategies, it can destigmatize women's health issues and increase support. More specifically, a targeted educational intervention will serve to increase social support and awareness in a cost-effective and sustainable manner, resulting in more open conversations and mitigation of complications, to improve the quality of life in menopausal women and help prepare younger women for their future menopausal journey.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12905-023-02641-4>.

Additional file 1. Participant demographic.

Additional file 2. All survey questions and corresponding answer options.

Acknowledgements

Not applicable.

Authors' contributions

VP designed and coordinated the study, summarized the data, and drafted the manuscript. BCS assisted in designing the study, analyzed the data, generated the figure and tables, and contributed to manuscript writing. SR contributed to conceptualizing the study, and edited the manuscript. All authors read and approved the final manuscript.

Funding

This research has been funded by the Women and Children's Health Research Institute through the generosity of the Alberta Women's Health Foundation and through generous support from the Cavarzan Chair in Mature Women's Health awarded to SR. VP was supported by a studentship from the Women and Children's Health Research Institute of the University of Alberta and by a Mach-Gaensslen Studentship award.

Availability of data and materials

The data pertaining to the current study are available from the corresponding authors in accordance with appropriate data use agreements.

Declarations

Ethics approval and consent to participate

The study received ethics approval from the University of Alberta Health Research Ethics Board (Pro00108310). Informed consent was obtained from all participants. All the methods included in this study are in accordance with the declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Obstetrics and Gynecology, Faculty of Medicine & Dentistry, Women and Children's Health Research Institute, University of Alberta, 626-1 Community Service Centre, Royal Alexandra Hospital, 10240 Kingsway Ave, Edmonton T5H-3V9, Canada.

Received: 4 April 2023 Accepted: 7 September 2023

Published online: 15 September 2023

References

- Davis SR, Lambrinoukaki I, Lumsden M, Mishra GD, Pal L, Rees M, et al. Menopause Nat Rev Dis Primers. 2015;1:15004. <https://doi.org/10.1038/nrdp.2015.4>.
- Yoshany N, Morowatisharifabad MA, Mihanpour H, Bahri N, Jadgal KM. The Effect of Husbands' Education Regarding Menopausal Health on Marital Satisfaction of Their Wives. *J Menopausal Med*. 2017;23(1):15–24. <https://doi.org/10.6118/jmm.2017.23.1.15>.
- Polat F, Orhan I, Küçükkeleşçe DŞ. Does social support affect menopausal symptoms in menopausal women? *Perspect Psychiatr Care*. 2022;58(3):1062–70. <https://doi.org/10.1111/ppc.12899>.
- Maharjan RK. Impact of educational intervention on reduction of menopausal effects. *J Patan Acad Health Sci*. 2014;1(2):55–7. <https://doi.org/10.3126/jpahs.v1i2.20183>.
- Namazi M, Sadeghi R, Behboodi ZM. Social Determinants of Health in Menopause: An Integrative Review. *Int J Womens Health*. 2019;11:637–47. <https://doi.org/10.2147/IJWH.S228594>.
- Bromberger JT, Kravitz HM. Mood and Menopause: Findings from the Study of Women's Health Across the Nation (SWAN) over ten years. *Obstet Gynecol Clin North Am*. 2011;38(3):609–25. <https://doi.org/10.1016/j.ogc.2011.05.011>.
- Chedraui P, San Miguel G, Avila C. Quality of life impairment during the female menopause transition is related to personal and partner factors. *Gynecol Endocrinol*. 2009;25(2):130–5. <https://doi.org/10.1080/09513590802617770>.
- Sydora BC, Alvdaj T, Malley A, Mayan M, Shandro T, Ross S. Walking together: women with the severe symptoms of menopause propose a platform for a walking program; outcome from focus groups. *BMC Womens Health*. 2020;20(1):165. <https://doi.org/10.1186/s12905-020-01037-y>.
- Sydora BC, Graham B, Oster RT, Ross S. Menopause experience in First Nations women and initiatives for menopause symptom awareness; a community-based participatory research approach. *BMC Womens Health*. 2021;21(1):179. <https://doi.org/10.1186/s12905-021-01303-7>.
- Shariat SM, Simbar M, Dolatian M, Nasiri M. The relationship between perceived social support and women experiences in menopause. *Advances in Nursing & Midwifery*. 2016;25(90):55–64 (<https://journals.sbm.u.ac.ir/en-jnm/article/view/11644>).
- Bahri N, Yoshany N, Morowatisharifabad MA, Noghabi AD, Sajjadi M. The effects of menopausal health training for spouses on women's quality of life during menopause transitional period. *Menopause*. 2016;23(2):183–8. <https://doi.org/10.1097/GME.0000000000000588>.
- Sharma A, Minh Duc NT, Luu Lam Thang T, Nam NH, Ng SJ, Abbas KS, et al. A Consensus-Based Checklist for Reporting of Survey Studies (CROSS). *J Gen Intern Med*. 2021;36(10):3179–87.
- Eysenbach G. Improving the Quality of Web Surveys: The Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res*. 2004;6(3):e34. <https://doi.org/10.2196/jmir.6.3.e34>. (PMID: 15471760).
- Baber RJ, Panay N, Fenton A, IMS Writing Group. 2016 IMS Recommendations on women's midlife health and menopause hormone therapy. *Climacteric*. 2016;19(2):109–50. <https://doi.org/10.3109/13697137.2015.1129166>.
- North American Menopause Society (NAMS). *The Menopause Guidebook*, 9th edition. 2020.
- Heinemann K, Ruebig A, Potthoff P, Schneider HP, Strelow F, Heinemann LA, Do MT. The Menopause Rating Scale (MRS) scale: a methodological review. *Health Qual Life Outcomes*. 2004;2:45. <https://doi.org/10.1186/1477-7525-2-45>.
- Hilditch JR, Lewis J, Peter A, van Maris B, Ross A, Franssen E, Guyatt GH, Norton PG, Dunn E. A menopause-specific quality of life questionnaire: development and psychometric properties. *Maturitas*. 1996;24(3):161–75. [https://doi.org/10.1016/s0378-5122\(96\)82006-8](https://doi.org/10.1016/s0378-5122(96)82006-8).
- Lewis JE, Hilditch JR, Wong CJ. Further psychometric property development of the Menopause-Specific Quality of Life questionnaire and development of a modified version MENQOL-Intervention questionnaire. *Maturitas*. 2005;50(3):209–21. <https://doi.org/10.1016/j.maturitas.2004.06.015>.
- Sydora BC, Fast H, Campbell S, Yuksel N, Lewis JE, Ross S. Use of the Menopause-Specific Quality of Life (MENQOL) questionnaire in research and clinical practice: a comprehensive scoping review. *Menopause*. 2016;23:1038–51. <https://doi.org/10.1097/GME.0000000000000636>.
- Sydora BC, Yuksel N, Veltri NL, et al. Patient characteristics, menopause symptoms, and care provided at an interdisciplinary menopause clinic: retrospective chart review. *Menopause*. 2017;25:102–5. <https://doi.org/10.1097/GME.0000000000000942>.
- Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377–81. <https://doi.org/10.1016/j.jbi.2008.08.010>.
- Shahzad D, Thakur AA, Kidwai S, Shaikh HO, AlSuwaidi AO, AlOtaifi AF, et al. Women's knowledge and awareness on menopause symptoms and its treatment options remains inadequate: a report from the United Arab Emirates. *Menopause*. 2021;28(8):918–27. <https://doi.org/10.1097/GME.0000000000001783>.
- Parish SJ, Faubion SS, Weinberg M, Bernick B, Mirkin S. The MATE survey: men's perceptions and attitudes towards menopause and their role in partners' menopausal transition. *Menopause*. 2019;26(10):1110–6. <https://doi.org/10.1097/GME.0000000000001373>.
- Beadle JN, Paradiso S, Kovach C, Polgreen L, Denburg NL, Tranel D. Effects of age-related differences in empathy on social economic decision-making. *Int Psychogeriatr*. 2012;24(5):822–33.
- Andrews NCZ, McDowell H, Spadafora N, Andrews DV. Using social network position to understand early adolescents' power and dominance within a school context. *Sch Psychol*. 2022;37(6):445–54. <https://doi.org/10.1037/spq0000445>.
- Shariat SM, Ebrahimzadeh M, Safari A, Jamali J. The Effectiveness of Husbands' Awareness of Menopausal Complications on Physical Psychological Experiences of Postmenopausal Women. *J Holist Nurs*. 2021;31:155–64. <https://doi.org/10.32598/jhnm.31.3.2070>.
- Currie H, Moger SJ. Menopause - Understanding the impact on women and their partners. *Post Reprod Health*. 2019;25(4):183–90. <https://doi.org/10.1177/2053369119895413>.
- Çaçapava JR, Cid B, Komura LA, Lima P. Men's perceptions and attitudes toward their wives experiencing menopause. *J Women Aging*. 2016;28(4):322–33. <https://doi.org/10.1080/08952841.2015.1017430>.
- Harper JC, Phillips S, Biswakarma R, Yasmin E, Saridogan E, Radhakrishnan S, et al. An online survey of perimenopausal women to determine their attitudes and knowledge of the menopause. *Womens Health (Lond)*. 2022;18:17455057221106890. <https://doi.org/10.1177/17455057221106890>.
- Baquedano L, Fasero M, Gabasa L, Coronado P, Presa J, Mendoza N. What do Spanish women know about menopause? COMEM study. *J Obstet Gynaecol*. 2022;42(5):1448–54. <https://doi.org/10.1080/01443615.2021.1998892>.

31. Koyuncu T, Unsal A, Arslantas D. Evaluation of the Effectiveness of Health Education on Menopause Symptoms and Knowledge and Attitude in Terms of Menopause. *J Epidemiol Glob Health*. 2018;8(1–2):8–12. <https://doi.org/10.2991/jjegh.2018.08.103>.
32. Woods NF, Mitchell ES. Symptoms during the perimenopause: prevalence, severity, trajectory, and significance in women's lives. *Am J Med*. 2005;118(Suppl 12B):14–24. <https://doi.org/10.1016/j.amjmed.2005.09.031>.
33. Maki PM, Kornstein SG, Joffe H, Bromberger JT, Freeman EW, Athappilly G, Bobo WV, Rubin LH, Koleva HK, Cohen LS, Soares CN, Board of Trustees for The North American Menopause Society (NAMS) and the Women and Mood Disorders Task Force of the National Network of Depression Centers. Guidelines for the evaluation and treatment of perimenopausal depression: summary and recommendations. *Menopause*. 2018;25(10):1069–85. <https://doi.org/10.1097/GME.0000000000001174>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

