


CORRECTION

Open Access



Correction to: Sociodemographic correlates of colorectal cancer screening completion among women adherent to mammography screening guidelines by place of birth

Deeonna E. Farr^{1*} , Leslie E. Cofe¹ , Alison T. Brenner^{2,4} , Ronny A. Bell³  and Daniel S. Reuland^{2,4} 

Correction to: BMC Women's Health (2022) 22:125
<https://doi.org/10.1186/s12905-022-01694-1>

The original article has been corrected.

Following publication of the original article [1], there is formatting error in Tables 1 and 2. The Tables 1 and Table 2 should be as shown below:

The original article can be found online at <https://doi.org/10.1186/s12905-022-01694-1>.

*Correspondence: farrd17@ecu.edu

¹ Department of Health Education and Promotion, College of Health and Human Performance, East Carolina University, 2307 Carol G. Belk Building, Mail Stop 529, Greenville, NC 27858, USA
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Table 1 Sociodemographic characteristics of women adherent to breast cancer guidelines, by place of birth, NHIS 2015

	Total	Place of birth n (%)		p value
		US-born	Foreign-born	
CRC screening adherent				
Yes	665 (55.52%)	515 (58.26%)	150 (46.46%)	0.0005
No	541 (44.48%)	374 (41.74%)	167 (53.54%)	
Types of CRC screening				
Sigmoidoscopy				
Yes	16 (1.47%)	9 (1.27%)	7 (2.13%)	0.2910
No	1182 (98.53%)	875 (98.73%)	307 (97.87%)	
Colonoscopy				
Yes	602 (50.33%)	470 (52.77%)	132 (42.21%)	0.0015
No	595 (49.67%)	414 (47.23%)	181 (57.49%)	
Fecal occult blood test				
Yes	124 (10.81%)	94 (11.31%)	30 (9.16%)	0.2901
No	1078 (89.19%)	793 (88.69%)	285 (90.84%)	
Personal characteristics				
Age				
50–64	841 (70.98%)	630 (72.37%)	211 (66.39%)	0.056
65+	365 (29.02%)	259 (27.63%)	106 (33.61%)	
Race/ethnicity				
White	607 (58.38%)	571 (70.80%)	36 (17.22%)	< 0.001
Hispanic	245 (16.14%)	65 (5.56%)	180 (51.13%)	
Black	248 (18.03%)	220 (20.76%)	28 (8.99%)	
Asian	106 (7.48%)	33 (2.88%)	73 (22.67%)	
Degree				
No high school degree	193 (12.82%)	67 (6.24%)	126 (34.59%)	< 0.001
High school degree	289 (22.90%)	218 (23.42%)	71 (21.19%)	
Some college/associate degree	364 (30.80%)	311 (34.81%)	53 (17.53%)	
College degree or higher	358 (33.48%)	292 (35.53%)	66 (26.69%)	
Federal poverty level				
≤ 138%	280 (19.48%)	168 (16.22%)	112 (30.27%)	< 0.001
139–200%	126 (9.38%)	83 (8.28%)	43 (13.02%)	
210–400%	328 (26.95%)	243 (27.55%)	85 (24.96%)	
≥ 410%	472 (44.19%)	395 (47.95%)	77 (31.75%)	
Marital status				
Married	593 (53.56%)	428 (53.30%)	165 (54.42%)	0.411
Widowed/divorced/separated	469 (36.81%)	345 (36.51%)	124 (37.81%)	
Single	140 (9.63%)	112 (10.19%)	28 (7.78%)	
Region				
Northeast	218 (18.72%)	134 (15.75%)	84 (28.57%)	< 0.001
North Central/Midwest	204 (18.41%)	172 (20.39%)	32 (11.86%)	
South	439 (40.00%)	353 (43.38%)	86 (28.83%)	
West	345 (22.87%)	230 (20.48%)	115 (30.74%)	
Language spoken				
English	796 (69.65%)	749 (85.41%)	47 (17.54%)	< 0.001
Mostly English	154 (13.44%)	105 (11.75%)	49 (10.02%)	
Only Spanish/other language	137 (9.34%)	6 (0.65%)	131 (38.09%)	
Mostly Spanish	48 (3.05%)	0 (0.00%)	48 (13.14%)	
Spanish and English equally	71 (4.51%)	29 (2.19%)	42 (12.19%)	

Table 1 (continued)

	Total	Place of birth n (%)		p value
		US-born	Foreign-born	
Place of birth				
Foreign-born	317 (23.22%)	–	–	–
U.S.-born	889 (76.78%)	–	–	–
Insurance				
Private	614 (55.96%)	468 (57.55%)	146 (50.73%)	<0.001
Medicaid	111 (7.33%)	70 (6.65%)	41 (9.55%)	
Medicare	158 (12.24%)	109 (11.10%)	49 (16.01%)	
Dual eligible	62 (4.01%)	33 (2.84%)	29 (8.24%)	
Other	199 (16.61%)	172 (18.53%)	27 (8.90%)	
None	59 (3.76%)	34 (2.91%)	25 (6.57%)	
Breast cancer risk				
More likely to get cancer	48 (4.72%)	33 (4.35%)	15 (5.67%)	0.460
Less likely	589 (49.80%)	445 (50.47%)	144 (47.54%)	
About as likely	501 (45.49%)	364 (45.19%)	137 (46.50%)	
CRC risk				
More likely to get cancer	17 (1.03%)	7 (0.57%)	10 (2.56%)	0.008
Less likely	632 (56.55%)	479 (57.62%)	153 (52.95%)	
About as likely	475 (42.42%)	343 (41.80%)	132 (44.49%)	

Table 2 Factors associated with CRC screening adherence among women adherent to breast cancer screening, NHIS 2015

	CRC screening adherence			
	Unadjusted OR (95% CI)	Adjusted OR (95% CI) Overall	Adjusted OR (95% CI) US-born	Adjusted OR (95% CI) Foreign-born
Age				
50–64	Ref	Ref	Ref	Ref
65+	1.82 (1.34–2.47)	1.76 (1.06–2.91)	2.70 (1.37–5.34)	0.57 (0.28–1.16)
Race				
White	Ref	Ref	Ref	Ref
Hispanic	0.73 (0.55–0.98)	1.01 (0.59–1.72)	1.63 (0.79–3.38)	0.82 (0.27–2.50)
Black	0.94 (0.70–1.26)	0.95 (0.50–1.80)	1.10 (0.76–1.59)	3.69 (1.22–11.21)
Asian	0.68 (0.43–1.06)	1.28 (0.73–2.23)	0.65 (0.30–1.41)	1.05 (0.43–2.56)
Degree				
No high school degree	Ref	Ref	Ref	Ref
High school degree	1.21 (0.84–1.74)	1.07 (0.67–1.74)	1.13 (0.63–2.01)	1.23 (0.54–2.81)
Some college/associated degree	1.44 (1.01–2.06)	1.39 (0.87–2.23)	1.64 (0.92–2.91)	1.22 (0.45–3.34)
College degree	1.67 (1.16–2.34)	1.37 (0.79–2.35)	1.47 (0.76–2.83)	1.52 (0.58–4.03)
Federal poverty level				
≤ 138%	Ref	Ref	Ref	Ref
139–200%	0.91 (0.59–1.41)	0.80 (0.48–1.33)	0.77 (0.39–1.51)	0.92 (0.40–2.09)
210–400%	1.00 (0.7–1.44)	0.93 (0.60–1.46)	0.92 (0.52–1.64)	0.95 (0.47–1.93)
≥ 410%	1.50 (1.09–2.08)	1.43 (0.88–2.35)	1.55 (0.82–2.93)	1.13 (0.54–2.34)
Insurance				
Private	5.19 (2.44–11.03)	3.84 (1.83–8.09)	3.32 (1.33–8.27)	6.94 (1.42–33.88)
Medicaid	3.45 (1.53–7.77)	2.99 (1.33–6.76)	2.31 (0.82–6.49)	6.91 (1.36–35.13)
Medicare	7.39 (3.29–16.61)	4.15 (1.73–9.94)	2.95 (1.02–8.54)	18.65 (2.85–122.12)
Dual eligible	5.71 (2.15–15.20)	3.48 (1.17–10.33)	2.03 (0.48–8.54)	20.47 (3.36–124.96)
Other	9.16 (4.16–20.15)	5.11 (2.10–12.42)	2.95 (1.02–8.54)	41.51 (5.64–305.55)

Table 2 (continued)

	CRC screening adherence			
	Unadjusted OR (95% CI)	Adjusted OR (95% CI) Overall	Adjusted OR (95% CI) US-born	Adjusted OR (95% CI) Foreign-born
None	Ref	Ref	Ref	Ref
Marital status				
Married	Ref	Ref	Ref	Ref
Widowed/divorced/separated	1.08 (0.84–1.41)	1.08 (0.79–1.48)	1.20 (0.82–1.76)	0.97 (0.54–1.72)
Single	1.15 (0.75–1.77)	1.36 (0.83–2.23)	1.74 (0.97–3.10)	0.77 (0.31–1.96)
Region				
Northeast	Ref	Ref	Ref	Ref
North Central/Midwest	0.87 (0.55–1.35)	1.01 (0.60–1.70)	1.24 (0.70–2.20)	0.83 (0.33–2.09)
South	0.86 (0.58–1.27)	0.80 (0.52–1.22)	0.82 (0.51–1.320)	0.68 (0.27–1.74)
West	0.86 (0.58–1.28)	0.89 (0.56–1.41)	0.68 (0.41–1.12)	1.38 (0.58–3.28)
Language spoken				
English	Ref	Ref	Ref	Ref
Mostly English	1.07 (0.77–1.50)	1.04 (0.62–1.74)	1.07 (0.73–1.59)	0.91 (0.33–2.46)
Only Spanish/other language	0.55 (0.38–0.80)	1.81 (0.53–1.23)	0.26 (0.04–1.57)	1.37 (0.57–3.29)
Mostly Spanish	0.84 (0.47–1.51)	0.91 (0.57–1.44)	–	3.11 (1.03–9.42)
Spanish and English equally	1.16 (0.73–1.83)	3.84 (1.83–8.09)	–	3.03 (1.08–8.54)
Place of birth				
Foreign-born	0.62 (0.48–0.81)	0.53 (0.34–0.83)	–	–
U.S.-born	Ref	Ref	–	–
Breast cancer risk				
More likely to get cancer	Ref	Ref	Ref	Ref
Less likely	0.72 (0.40–1.32)	0.56 (0.27–1.16)	0.50 (0.21–1.18)	0.72 (0.16–3.26)
About as likely	0.61 (0.33–1.11)	0.56 (0.27–1.16)	0.52 (0.22–1.23)	0.48 (0.13–1.78)
CRC risk				
More likely to get cancer	1.68 (0.61–4.62)	Ref	Ref	Ref
Less likely	1.41 (0.51–3.89)	2.01 (0.70–5.73)	2.70 (0.59–12.31)	1.42 (0.35–5.84)
About as likely	Ref	1.82 (0.62–5.32)	2.05 (0.43–9.80)	2.16 (0.60–7.86)

Bolded AORs (95% CI) represent significant findings. All analyses were weighted to account for sampling

Author details

¹Department of Health Education and Promotion, College of Health and Human Performance, East Carolina University, 2307 Carol G. Belk Building, Mail Stop 529, Greenville, NC 27858, USA. ²Department of Medicine, Department of General Medicine and Clinical Epidemiology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA. ³Department of Social Sciences and Health Policy, Division of Public Health Sciences, Wake Forest School of Medicine, Medical Center Boulevard, Winston-Salem, NC 27157, USA. ⁴Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA.

Accepted: 13 July 2022

Published online: 01 August 2022

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.